

FIRST QUANTUM CORRECTION FOR THE MODULI SPACE OF STABLE BUNDLES OVER A RIEMANN SURFACE

VICENTE MUÑOZ

November, 1997

ABSTRACT. We compute some Gromov-Witten invariants of the moduli space M_Σ of odd degree rank two stable vector bundles over a Riemann surface Σ of genus $g \geq 2$. We thus find the first correction term for the quantum product of M_Σ and hence get the two leading terms of the relations satisfied by the natural generators of the quantum cohomology of M_Σ . Finally, we use this information to get a full description of the quantum cohomology of M_Σ when the genus of Σ is $g = 3$.

1. INTRODUCTION

Let Σ be a Riemann surface of genus $g \geq 2$ and let M_Σ denote the moduli space of flat $SO(3)$ -connections with nontrivial second Stiefel-Whitney class w_2 .

This is a smooth symplectic manifold of dimension $6g - 6$. Alternatively, we can consider Σ as a smooth complex curve of genus g . Fix a line bundle Λ on Σ of degree 1, then M_Σ is the moduli space of rank two stable vector bundles on Σ with determinant Λ , which is a smooth complex variety of complex dimension $3g - 3$. The symplectic deformation class of M_Σ only depends on g and not on the particular complex structure on Σ .

The manifold $X = M_\Sigma$ is a positive symplectic manifold with $\pi_2(X) = \mathbb{Z}$. For such a manifold X , its quantum cohomology, $QH^*(X)$, is well-defined (see [14] [15] [8] [12]). As vector spaces, $QH^*(X) = H^*(X)$ (rational coefficients are understood), but the multiplicative structure is different. Let A denote the positive generator of $\pi_2(X)$, i.e. the generator such that the symplectic form evaluated on A is positive. Let $N = c_1(X)[A] \in \mathbb{Z}_{>0}$. Then there is a natural $\mathbb{Z}/2N\mathbb{Z}$ -grading for $QH^*(X)$, which comes from reducing the $\mathbb{Z}$ -grading of $H^*(X)$. (For the case $X = M_\Sigma$, $N = 2$, so $QH^*(M_\Sigma)$ is $\mathbb{Z}/4\mathbb{Z}$ -graded). The ring structure of $QH^*(X)$, called quantum multiplication, is a deformation of the usual cup product for $H^*(X)$. For $\alpha \in H^p(X)$, $\beta \in H^q(X)$, we

*Supported by a grant from Ministerio de Educación y Cultura

1991 Mathematics Subject Classification. Primary: 57R57. Secondary: 58D27, 53C23, 57R57.

define the quantum product of α and β as

$$\alpha \cdot \beta = \sum_{d \geq 0} \Phi_{dA}(\alpha, \beta),$$

where $\Phi_{dA}(\alpha, \beta) \in H^{p+q-2Nd}(X)$ is given by $\langle \Phi_{dA}(\alpha, \beta), \gamma \rangle = \Psi_{dA}^X(\alpha, \beta, \gamma)$, the Gromov-Witten invariant, for all $\gamma \in H^{\dim X - p - q + 2Nd}(X)$. One has $\Phi_0(\alpha, \beta) = \alpha \cup \beta$. The other terms are the correction terms and all live in lower degree parts of the cohomology groups. It is a fact [15] that the quantum product gives an associative and graded commutative ring structure.

To define the Gromov-Witten invariant, let J be a generic almost complex structure compatible with the symplectic form. Then for every 2-homology class dA , $d \in \mathbb{Z}$, there is a moduli space $\mathcal{M}_{dA}$ of pseudoholomorphic rational curves (with respect to J) $f : \mathbb{P}^1 \rightarrow X$ with $f_*[\mathbb{P}^1] = dA$. Note that $\mathcal{M}_0 = X$ and that $\mathcal{M}_{dA}$ is empty for $d < 0$. For $d \geq 0$, the dimension of $\mathcal{M}_{dA}$ is $\dim X + 2Nd$. This moduli space $\mathcal{M}_{dA}$ admits a natural compactification, $\overline{\mathcal{M}}_{dA}$, called the Gromov-Uhlenbeck compactification [14] [15, section 3]. Consider now $r \geq 3$ different points $P_1, \dots, P_r \in \mathbb{P}^1$. Then we have defined an evaluation map $ev : \mathcal{M}_{dA} \rightarrow X^r$ by $f \mapsto (f(P_1), \dots, f(P_r))$. This map extends to $\overline{\mathcal{M}}_{dA}$ and its image, $ev(\overline{\mathcal{M}}_{dA})$, is a pseudo-cycle [15]. So for $\alpha_i \in H^{p_i}(M_\Sigma)$, $1 \leq i \leq r$, with $p_1 + \dots + p_r = \dim X + 2Nd$, we choose generic cycles A_i , $1 \leq i \leq r$, representatives of their Poincaré duals, and set

$$(1) \Psi_{dA}^X(\alpha_1, \dots, \alpha_r) = \langle A_1 \times \dots \times A_r, [ev(\overline{\mathcal{M}}_{dA})] \rangle = \# ev_{P_1}^*(A_1) \cap \dots \cap ev_{P_r}^*(A_r),$$

where $\#$ denotes count of points (with signs) and $ev_{P_i} : \mathcal{M}_{dA} \rightarrow X$, $f \mapsto f(P_i)$. This is a well-defined number and independent of the particular cycles. Also, as the manifold X is positive, $\text{coker } L_f = H^1(\mathbb{P}^1, f^*c_1(X)) = 0$, for all $f \in \mathcal{M}_{dA}$. By [14] the complex structure of X is generic and we can use it to compute the Gromov-Witten invariants.

Also for $r \geq 2$, let $\alpha_i \in H^{p_i}(M_\Sigma)$, $1 \leq i \leq r$, then

$$\alpha_1 \cdots \alpha_r = \sum_{d \geq 0} \Phi_{dA}(\alpha_1, \dots, \alpha_r),$$

where the correction terms $\Phi_{dA}(\alpha_1, \dots, \alpha_r) \in H^{p_1 + \dots + p_r - 2Nd}(X)$ are determined by $\langle \Phi_{dA}(\alpha_1, \dots, \alpha_r), \gamma \rangle = \Psi_{dA}^X(\alpha_1, \dots, \alpha_r, \gamma)$, for any $\gamma \in H^{\dim X + 2Nd - (p_1 + \dots + p_r)}(X)$.

Returning to our manifold $X = M_\Sigma$, there is a classical conjecture relating the quantum cohomology $QH^*(M_\Sigma)$ and the instanton Floer cohomology of $\Sigma \times \mathbb{S}^1$, $HF^*(\Sigma \times \mathbb{S}^1)$ (see [10]). In [1] a presentation of $QH^*(M_\Sigma)$ was given using physical methods, and in [10] it was proved that such a presentation was a presentation of $HF^*(\Sigma \times \mathbb{S}^1)$ indeed. Siebert and Tian have a program [16] to find the presentation of $QH^*(M_\Sigma)$, which goes through proving a recursion formula for the Gromov-Witten invariants of M_Σ in terms of the genus g . This will complete the proof of the conjectural isomorphism $QH^*(M_\Sigma) \cong HF^*(\Sigma \times \mathbb{S}^1)$.

The purpose of this paper is two-fold. On the one hand we aim to compute the Gromov-Witten invariants $\Psi_A^{M_\Sigma}$, relating them to the Donaldson invariants for the algebraic surface $S = \Sigma \times \mathbb{P}^1$. This gives the first correction term of the quantum product of M_Σ and hence the first two leading terms of the relations satisfied by the generators of $QH^*(M_\Sigma)$. In particular we obtain the coefficient c_g of [16, section 3.3]. On the other hand we infer a full presentation of $QH^*(M_\Sigma)$ when the genus of Σ is $g = 3$ (recall that the case of genus $g = 2$ was worked out by Donaldson [3]). This is the starting point of the induction in [16].

The paper is organised as follows. In section 2 we review the ordinary cohomology ring of M_Σ . In section 3 the moduli space of lines in M_Σ is described, in order to compute the corresponding Gromov-Witten invariants in section 4. In section 5 we study the quantum cohomology of M_Σ and finally determine it completely in the case of genus $g = 3$ in section 6.

Acknowledgements: I want to thank my D. Phil. supervisor Simon Donaldson, for his encouragement and invaluable help. Also I am very grateful to Bernd Siebert and Gang Tian for helpful conversations and for letting me have a copy of their preprint [16].

2. CLASSICAL COHOMOLOGY RING OF M_Σ

Let us recall the known description of the homology of M_Σ [6] [17] [10]. Let $\mathcal{U} \rightarrow \Sigma \times M_\Sigma$ be the universal bundle and consider the Künneth decomposition of

$$(2) \quad c_2(\text{End}_0 \mathcal{U}) = 2[\Sigma] \otimes \alpha + 4\psi - \beta,$$

with $\psi = \sum \gamma_i \otimes \psi_i$, where $\{\gamma_1, \dots, \gamma_{2g}\}$ is a symplectic basis of $H^1(\Sigma; \mathbb{Z})$ with $\gamma_i \gamma_{i+g} = [\Sigma]$ for $1 \leq i \leq g$ (also $\{\gamma_i^\#\}$ will denote the dual basis for $H_1(\Sigma; \mathbb{Z})$). Here we can suppose without loss of generality that $c_1(\mathcal{U}) = \Lambda + \alpha$ (see [17]). In terms of the map $\mu : H_*(\Sigma) \rightarrow H^{4-*}(M_\Sigma)$, given by $\mu(a) = -\frac{1}{4} p_1(\mathfrak{g}_\mathcal{U})/a$ (here $\mathfrak{g}_\mathcal{U} \rightarrow \Sigma \times M_\Sigma$ is the associated universal $SO(3)$ -bundle, and $p_1(\mathfrak{g}_\mathcal{U}) \in H^4(\Sigma \times M_\Sigma)$ its first Pontrjagin class), we have

$$\begin{cases} \alpha = 2\mu(\Sigma) \in H^2 \\ \psi_i = \mu(\gamma_i^\#) \in H^3, \\ \beta = -4\mu(x) \in H^4 \end{cases} \quad 1 \leq i \leq 2g$$

where $x \in H_0(\Sigma)$ is the class of the point, and $H^i = H^i(M_\Sigma)$. These elements generate $H^*(M_\Sigma)$ as a ring [6] [19], and α is the positive generator of $H^2(M_\Sigma; \mathbb{Z})$. We can rephrase this as saying that there exists an epimorphism

$$(3) \quad \mathbb{A}(\Sigma) = \mathbb{Q}[\alpha, \beta] \otimes \Lambda(\psi_1, \dots, \psi_{2g}) \twoheadrightarrow H^*(M_\Sigma)$$

(the notation $\mathbb{A}(\Sigma)$ follows that of Kronheimer and Mrowka [7]). The mapping class group $\text{Diff}(\Sigma)$ acts on $H^*(M_\Sigma)$, with the action factoring through the action of $\text{Sp}(2g, \mathbb{Z})$ on $\{\psi_i\}$. The invariant part, $H_I^*(M_\Sigma)$, is generated by α , β and $\gamma = -2 \sum_{i=0}^g \psi_i \psi_{i+g}$. Then there is an epimorphism

$$(4) \quad \mathbb{Q}[\alpha, \beta, \gamma] \twoheadrightarrow H_I^*(M_\Sigma)$$

which allows us to write

$$H_I^*(M_\Sigma) = \mathbb{Q}[\alpha, \beta, \gamma]/I_g,$$

where I_g is the ideal of relations satisfied by α , β and γ . Recall that $\deg(\alpha) = 2$, $\deg(\beta) = 4$ and $\deg(\gamma) = 6$. From [17], a basis for $H_I^*(M_\Sigma)$ is given by the monomials $\alpha^a \beta^b \gamma^c$, with $a + b + c < g$. For $0 \leq k \leq g$, the primitive component of $\Lambda^k H^3$ is

$$\Lambda_0^k H^3 = \ker(\gamma^{g-k+1} : \Lambda^k H^3 \rightarrow \Lambda^{2g-k+2} H^3).$$

Then the $\text{Sp}(2g, \mathbb{Z})$ -decomposition of $H^*(M_\Sigma)$ is [6]

$$H^*(M_\Sigma) = \bigoplus_{k=0}^g \Lambda_0^k H^3 \otimes \mathbb{Q}[\alpha, \beta, \gamma]/I_{g-k}.$$

Proposition 1 ([17]). *For $g = 1$, let $q_1^1 = \alpha$, $q_1^2 = \beta$, $q_1^3 = \gamma$. Define recursively, for $g \geq 1$,*

$$\begin{cases} q_{g+1}^1 = \alpha q_g^1 + g^2 q_g^2 \\ q_{g+1}^2 = \beta q_g^1 + \frac{2g}{g+1} q_g^3 \\ q_{g+1}^3 = \gamma q_g^1 \end{cases}$$

Then $I_g = (q_g^1, q_g^2, q_g^3)$, for all $g \geq 1$. Note that $\deg(q_g^1) = 2g$, $\deg(q_g^2) = 2g + 2$ and $\deg(q_g^3) = 2g + 4$.

3. HOLOMORPHIC LINES IN M_Σ

In order to compute the Gromov-Witten invariants $\Psi_A^{M_\Sigma}$, we need to describe the space of lines, i.e. rational curves in M_Σ representing the generator $A \in H_2(M_\Sigma; \mathbb{Z})$,

$$\mathcal{M}_A = \{f : \mathbb{P}^1 \rightarrow M_\Sigma/f \text{ holomorphic, } f_*[\mathbb{P}^1] = A\}.$$

Let us fix some notation. Let J denote the Jacobian variety of Σ parametrising line bundles of degree 0 and let $\mathcal{L} \rightarrow \Sigma \times J$ be the universal line bundle. If $\{\gamma_i\}$ is the basis of $H^1(\Sigma)$ introduced in section 2 then $c_1(\mathcal{L}) = \sum \gamma_i \otimes \phi_i \in H^1(\Sigma) \otimes H^1(J)$, where $\{\phi_i\}$ is a symplectic basis for $H^1(J)$. Thus $c_1(\mathcal{L})^2 = -2[\Sigma] \otimes \omega \in H^2(\Sigma) \otimes H^2(J)$, where $\omega = \sum_{i=1}^g \phi_i \wedge \phi_{i+g}$ is the natural symplectic form for J .

Consider now the algebraic surface $S = \Sigma \times \mathbb{P}^1$. It has irregularity $q = g \geq 2$, geometric genus $p_g = 0$ and canonical bundle $K \equiv -2\Sigma + (2g - 2)\mathbb{P}^1$. Recall that Λ is a fixed line bundle of degree 1 on Σ . Fix the line bundle $L = \Lambda \otimes \mathcal{O}_{\mathbb{P}^1}(1)$ on S (we omit all pull-backs) with $c_1 = c_1(L) \equiv \mathbb{P}^1 + \Sigma$, and put $c_2 = 1$. The ample cone of S

is $\{a\mathbb{P}^1 + b\Sigma / a, b > 0\}$. Let H_0 be a polarisation close to $\mathbb{P}^1$ in the ample cone and H be a polarisation close to Σ , i.e. $H = \Sigma + t\mathbb{P}^1$ with t small. We wish to study the moduli space $\mathfrak{M} = \mathfrak{M}_H(c_1, c_2)$ of H -stable bundles over S with Chern classes c_1 and c_2 .

Proposition 2. *$\mathfrak{M}$ can be described as a bundle $\mathbb{P}^{2g-1} \rightarrow \mathfrak{M} = \mathbb{P}(\mathcal{E}_\zeta^\vee) \rightarrow J$, where $\mathcal{E}_\zeta$ is a bundle on J with $\text{ch } \mathcal{E}_\zeta = 2g + 8\omega$. So $\mathfrak{M}$ is compact, smooth and of the expected dimension $6g - 2$. The universal bundle $\mathcal{V} \rightarrow S \times \mathfrak{M}$ is given by*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^1}(1) \otimes \mathcal{L} \otimes \lambda \rightarrow \mathcal{V} \rightarrow \Lambda \otimes \mathcal{L}^{-1} \rightarrow 0,$$

where λ is the tautological line bundle for $\mathfrak{M}$.

Proof. For the polarisation H_0 , the moduli space of H_0 -stable bundles with Chern classes c_1, c_2 is empty by [13]. Now for $p_1 = -4c_2 + c_1^2 = -2$ there is only one wall, determined by $\zeta \equiv -\mathbb{P}^1 + \Sigma$ (here we fix $\zeta = 2\Sigma - L = \Sigma - c_1(\Lambda)$ as a divisor), so the moduli space of H -stable bundles with Chern classes c_1, c_2 is obtained by crossing the wall as described in [9]. First, note that the results in [9] use the hypothesis of $-K$ being effective, but the arguments work equally well with the weaker assumption of ζ being a good wall [9, remark 1] (see also [5] for the case of $q = 0$). In our case, $\zeta \equiv -\mathbb{P}^1 + \Sigma$ is a good wall (i.e. $\pm\zeta + K$ are both not effective) with $l_\zeta = 0$. Now with the notations of [9], F is a divisor such that $2F - L \equiv \zeta$, i.e. $F = \Sigma$. Also $\mathcal{F} \rightarrow S \times J$ is the universal bundle parametrising divisors homologically equivalent to F , i.e. $\mathcal{F} = \mathcal{L} \otimes \mathcal{O}_{\mathbb{P}^1}(1)$. Let $\pi : S \times J \rightarrow J$ be the projection. Then $\mathfrak{M} = E_\zeta = \mathbb{P}(\mathcal{E}_\zeta^\vee)$, where

$$\mathcal{E}_\zeta = \mathcal{E}\text{xt}_\pi^1(\mathcal{O}(L - \mathcal{F}), \mathcal{O}(\mathcal{F})) = R^1\pi_*(\mathcal{O}(\zeta) \otimes \mathcal{L}^2).$$

Actually $\mathfrak{M}$ is exactly the set of bundles E that can be written as extensions

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^1}(1) \otimes L \rightarrow E \rightarrow \Lambda \otimes L^{-1} \rightarrow 0$$

for a line bundle L of degree 0. The Chern character is [9, section 3] $\text{ch } \mathcal{E}_\zeta = 2g + e_{K-2\zeta}$, where $e_\alpha = -2(\mathbb{P}^1 \cdot \alpha)\omega$ (the class Σ defined in [9, lemma 11] is $\mathbb{P}^1$ in our case). Finally, the description of the universal bundle follows from [9, theorem 10]. $\square$

Proposition 3. *There is a well defined map $\mathcal{M}_A \rightarrow \mathfrak{M}$.*

Proof. Every line $f : \mathbb{P}^1 \rightarrow M_\Sigma$ gives a bundle $E = (\text{id}_\Sigma \times f)^*\mathcal{U}$ over $\Sigma \times \mathbb{P}^1$ by pulling-back the universal bundle $\mathcal{U} \rightarrow \Sigma \times M_\Sigma$. Then for any $t \in \mathbb{P}^1$, the bundle $E|_{\Sigma \times t}$ is defined by $f(t)$. Now, by equation (2), $p_1(E) = p_1(\mathcal{U})[\Sigma \times A] = -2\alpha[A] = -2$. Since $c_1(E) = (\text{id}_\Sigma \times f)^*c_1(\mathcal{U}) = \Lambda + \Sigma$, it must be $c_2 = 1$. To see that E is H -stable, consider any sub-line bundle $L \hookrightarrow E$ with $c_1(L) \equiv a\mathbb{P}^1 + b\Sigma$. Restricting to any $\Sigma \times t \subset \Sigma \times \mathbb{P}^1$ and using the stability of $E|_{\Sigma \times t}$, one gets $a \leq 0$. Then $c_1(L) \cdot \Sigma < \frac{c_1(E) \cdot \Sigma}{2}$, which yields the H -stability of E (recall that H is close to Σ). So $E \in \mathfrak{M}$. $\square$

Now define N as the set of extensions on Σ of the form

$$(5) \quad 0 \rightarrow L \rightarrow E \rightarrow \Lambda \otimes L^{-1} \rightarrow 0,$$

for L a line bundle of degree 0. Then the groups $\text{Ext}^1(\Lambda \otimes L^{-1}, L) = H^1(L^2 \otimes \Lambda^{-1}) = H^0(L^{-2} \otimes \Lambda \otimes K)$ are of constant dimension g . Moreover $H^0(L^2 \otimes \Lambda^{-1}) = 0$, so the moduli space N which parametrises extensions like (1) is given as $N = \mathbb{P}(\mathcal{E}^\vee)$, where $\mathcal{E} = \mathcal{E}\text{xt}_p^1(\Lambda \otimes \mathcal{L}^{-1}, \mathcal{L}) = R^1p_*(\mathcal{L}^2 \otimes \Lambda^{-1})$, $p : \Sigma \times J \rightarrow J$ the projection. Then we have a fibration $\mathbb{P}^{g-1} \rightarrow N = \mathbb{P}(\mathcal{E}^\vee) \rightarrow J$. The Chern character of $\mathcal{E}$ is

$$(6) \quad \begin{aligned} \text{ch}(\mathcal{E}) &= \text{ch}(R^1p_*(\mathcal{L}^2 \otimes \Lambda^{-1})) = -\text{ch}(p_!(\mathcal{L}^2 \otimes \Lambda^{-1})) = \\ &= -p_*((\text{ch } \mathcal{L})^2 (\text{ch } \Lambda)^{-1} \text{Todd } T_\Sigma) = \\ &= -p_*((1 + c_1(\mathcal{L}) + \tfrac{1}{2}c_1(\mathcal{L})^2)(1 - \Lambda)(1 - \tfrac{1}{2}K)) = \\ &= -p_*(1 - \tfrac{1}{2}K + 2c_1(\mathcal{L}) - 4\omega \otimes [\Sigma] - \Lambda) = g + 4\omega. \end{aligned}$$

It is easy to check that all the bundles in N are stable, so there is a well-defined map

$$i : N \rightarrow M_\Sigma.$$

Now we wish to construct the space of lines in N . Note that $\pi_2(N) = \pi_2(\mathbb{P}^{g-1}) = \mathbb{Z}$, as there are no rational curves in J . Let $L \in \pi_2(N)$ be the positive generator. We want to describe

$$\mathcal{N}_L = \{f : \mathbb{P}^1 \rightarrow N/f \text{ holomorphic, } f_*[\mathbb{P}^1] = L\}.$$

For the projective space $\mathbb{P}^n$, the space of lines H_1 is the set of algebraic maps $f : \mathbb{P}^1 \rightarrow \mathbb{P}^n$ of degree 1. Such an f has the form $f[x_0, x_1] = [x_0u_0 + x_1u_1]$, $[x_0, x_1] \in \mathbb{P}^1$, where u_0, u_1 are linearly independent vectors in $\mathbb{C}^{n+1}$. So

$$H_1 = \mathbb{P}(\{(u_0, u_1)/u_0, u_1 \text{ are linearly independent}\}) \subset \mathbb{P}((\mathbb{C}^{n+1} \oplus \mathbb{C}^{n+1})^\vee) = \mathbb{P}^{2n+1}.$$

The complement of H_1 is the image of $\mathbb{P}^n \times \mathbb{P}^1 \hookrightarrow \mathbb{P}^{2n+1}$, $([u], [x_0, x_1]) \mapsto [x_0u, x_1u]$, which is a smooth n -codimensional algebraic subvariety. Now $\mathcal{N}_L$ can be described as

$$(7) \quad \begin{array}{ccccc} H_1 & \rightarrow & \mathcal{N}_L & \rightarrow & J \\ \cap & & \cap & & \parallel \\ \mathbb{P}^{2g-1} & \rightarrow & \mathbb{P}((\mathcal{E} \oplus \mathcal{E})^\vee) & \rightarrow & J \end{array}$$

Remark 4. Note that $\mathcal{E}_\zeta = R^1\pi_*(\mathcal{O}(\zeta) \otimes \mathcal{L}^2) = R^1\pi_*(\mathcal{O}_{\mathbb{P}^1}(1) \otimes \mathcal{L}^2 \otimes \Lambda^{-1}) = H^0(\mathcal{O}_{\mathbb{P}^1}(1)) \otimes R^1p_*(\mathcal{L}^2 \otimes \Lambda^{-1}) = H^0(\mathcal{O}_{\mathbb{P}^1}(1)) \otimes \mathcal{E} \cong \mathcal{E} \oplus \mathcal{E}$. So $\mathfrak{M} = \mathbb{P}((\mathcal{E} \oplus \mathcal{E})^\vee)$, canonically.

Proposition 5. *The map $i : N \rightarrow M_\Sigma$ induces a map $i_* : \mathcal{N}_L \rightarrow \mathcal{M}_A$. The composition $\mathcal{N}_L \rightarrow \mathcal{M}_A \rightarrow \mathfrak{M}$ is the natural inclusion of (7).*

Proof. The first assertion is clear as i is a holomorphic map. For the second, consider the universal sheaf on $\Sigma \times N$,

$$(8) \quad 0 \rightarrow \mathcal{L} \otimes U \rightarrow \mathbb{E} \rightarrow \Lambda \otimes \mathcal{L}^{-1} \rightarrow 0,$$

where $U = \mathcal{O}_N(1)$ is the tautological bundle of the fibre bundle $\mathbb{P}^{g-1} \rightarrow N \rightarrow J$. Any element in $\mathcal{N}_L$ is a line $\mathbb{P}^1 \hookrightarrow N$, which must lie inside a single fibre $\mathbb{P}^{g-1}$. Restricting (8) to this line, we have an extension

$$0 \rightarrow L \otimes \mathcal{O}_{\mathbb{P}^1}(1) \rightarrow E \rightarrow \Lambda \otimes L^{-1} \rightarrow 0$$

on $S = \Sigma \times \mathbb{P}^1$, which is the image of the given element in $\mathfrak{M}$ (here L is the line bundle corresponding to the fibre in which $\mathbb{P}^1$ sits). Now it is easy to check that the map $\mathcal{N}_L \rightarrow \mathfrak{M}$ is the inclusion of (7). $\square$

Corollary 6. i_* is an isomorphism.

Proof. By proposition 5, i_* has to be an open immersion. The group $PGL(2, \mathbb{C})$ acts on both spaces $\mathcal{N}_L$ and $\mathcal{M}_A$, and i_* is equivariant. The quotient $\mathcal{N}_L/PGL(2, \mathbb{C})$ is compact, being a fibration over the Jacobian with all the fibres the Grassmannian $\text{Gr}(\mathbb{C}^2, \mathbb{C}^{g-1})$, hence irreducible. As a consequence i_* is an isomorphism. $\square$

Remark 7. Notice that the lines in M_Σ are all contained in the image of N , which is of dimension $4g-2$ against $6g-6 = \dim M_\Sigma$. They do not fill all of M_Σ as one would naively expect.

4. COMPUTATION OF $\Psi_A^{M_\Sigma}$

The manifold N is positive with $\pi_2(N) = \mathbb{Z}$ and $L \in \pi_2(N)$ is the positive generator. Under the map $i : N \rightarrow M_\Sigma$, we have $i_*L = A$. Now $\dim N = 4g-2$ and $c_1(N)[L] = c_1(\mathbb{P}^{g-1})[L] = g$. So quantum cohomology of N , $QH^*(N)$, is well-defined and $\mathbb{Z}/2g\mathbb{Z}$ -graded. From corollary 6, it is straightforward to prove

Lemma 8. For any $\alpha_i \in H^{p_i}(M_\Sigma)$, $1 \leq i \leq r$, such that $p_1 + \dots + p_r = 6g-2$, it is $\Psi_A^{M_\Sigma}(\alpha_1, \dots, \alpha_r) = \Psi_L^N(i^*\alpha_1, \dots, i^*\alpha_r)$. $\square$

It is therefore important to know the Gromov-Witten invariants of N , i.e. its quantum cohomology. From the universal bundle (8), we can read the first Pontrjagin class $p_1(\mathfrak{g}_\mathbb{E}) = -8[\Sigma] \otimes \omega + h^2 - 2[\Sigma] \otimes h + 4h \cdot c_1(\mathcal{L}) \in H^4(\Sigma \times N)$, where $h = c_1(U)$ is the hyperplane class. So on N we have

$$(9) \quad \begin{cases} \alpha = 2\mu(\Sigma) = 4\omega + h \\ \psi_i = \mu(\gamma_i^\#) = -h \cdot \phi_i \\ \beta = -4\mu(x) = h^2 \end{cases}$$

Let us remark that h^2 denotes ordinary cup product in $H^*(N)$, a fact which will prove useful later. Now let us compute the quantum cohomology ring of N . The cohomology of J is $H^*(J) = \Lambda H_1$, where $H_1 = H_1(\Sigma)$. Now the fibre bundle description $\mathbb{P}^{2g-1} \rightarrow N = \mathbb{P}(\mathcal{E}^\vee) \rightarrow J$ implies that the usual cohomology of N is $H^*(N) = \Lambda H_1[h]/< h^g + c_1 h^{g-1} + \dots + c_g = 0 >$, where $c_i = c_i(\mathcal{E}) = \frac{4^i}{i!} \omega^i$, from (6). As the quantum cohomology has the same generators as the usual cohomology and the relations are a deformation of the usual relations [18], it must be $h^g + c_1 h^{g-1} + \dots + c_g = r$ in $QH^*(N)$, with $r \in \mathbb{Q}$. As in [15, example 8.5], r can be computed to be 1. So

$$(10) \quad QH^*(N) = \Lambda H_1[h]/< h^g + c_1 h^{g-1} + \dots + c_g = 1 >.$$

Lemma 9. *For any $s \in H^{2g-2i}(J)$, $0 \leq i \leq g$, denote by $s \in H^{2g-2i}(N)$ its pull-back to N under the natural projection. Then the quantum product $h^{2g-1+i}s$ in $QH^*(N)$ has component in $H^{4g-2}(N)$ equal to $\frac{(-8)^i}{i!} \omega^i \wedge s$.*

Proof. First note that for $s_1, s_2 \in H^*(J)$ such that their cup product in J is $s_1 s_2 = 0$, then the quantum product $s_1 s_2 \in QH^*(N)$ vanishes. This is so since every rational line in N is contained in a fibre of $\mathbb{P}^{2g-1} \rightarrow J \rightarrow N$.

Next note that $h^{g-1+i}s$ has component in $H^{4g-2}(N)$ equal to $s_i(\mathcal{E}) \wedge s = \frac{(-4)^i}{i!} \omega^i \wedge s$. Then multiply the standard relation (9) by h^{g-1+i} and work by induction on i . For $i = 0$ we get $h^{2g-1}s = h^{g-1}s$ and the assertion is obvious. For $i > 0$,

$$h^{2g-1+i}s + h^{2g-2+i}c_1s + \dots + h^{2g-1}c_is = h^{g-1+i}s.$$

So the component of $h^{2g-1+i}s$ in $H^{4g-2}(N)$ is

$$-\sum_{j=1}^i \frac{(-8)^{i-j}}{(i-j)!} \omega^{i-j} c_j s + \frac{(-4)^i}{i!} \omega^i s = \frac{(-8)^i}{i!} \omega^i s - \sum_{j=0}^i \frac{(-8)^{i-j} 4^j}{(i-j)! j!} \omega^i s + \frac{(-4)^i}{i!} \omega^i s = \frac{(-8)^i}{i!} \omega^i s.$$

□

Lemma 10. *Suppose $g > 2$. Let $\alpha^a \beta^b \psi_{i_1} \dots \psi_{i_r} \in \mathbb{A}(\Sigma)$ have degree $6g - 2$. Then*

$$\Psi_L^N(\alpha, \cdot^{(a)}, \alpha, \beta, \cdot^{(b)}, \beta, \psi_{i_1}, \dots, \psi_{i_r}) = < (4\omega + X)^a (X^2)^b \phi_{i_1} \dots \phi_{i_r} X^r, [J] >,$$

evaluated on J , where $X^{2g-1+i} = \frac{(-8)^i}{i!} \omega^i \in H^(J)$.*

Proof. By definition the left hand side is the component in $H^{4g-2}(N)$ of the quantum product $\alpha^a \beta^b \psi_{i_1} \dots \psi_{i_r} \in QH^*(N)$. From (9), this quantum product is $(4\omega + h)^a (h^2)^b (-h\phi_{i_1}) \dots (-h\phi_{i_r})$, upon noting that when $g > 2$, $\beta = h^2$ as a quantum product as there are no quantum corrections because of the degree. Note that r is even, so the statement of the lemma follows from lemma 9. □

Now we are in the position of relating the Gromov-Witten invariants $\Psi_A^{M_\Sigma}$ with the Donaldson invariants for $S = \Sigma \times \mathbb{P}^1$ (for definition of Donaldson invariants see [4] [7]).

Theorem 11. Suppose $g > 2$. Let $\alpha^a \beta^b \psi_{i_1} \cdots \psi_{i_r} \in \mathbb{A}(\Sigma)$ have degree $6g - 2$. Then

$$\Psi_A^{M_\Sigma}(\alpha, \cdot^{(a)}, \alpha, \beta, \cdot^{(b)}, \beta, \psi_{i_1}, \dots, \psi_{i_r}) = (-1)^{g-1} D_{S,H}^{c_1}((2\Sigma)^a (-4pt)^b \gamma_{i_1}^\# \cdots \gamma_{i_r}^\#),$$

where $D_{S,H}^{c_1}$ stands for the Donaldson invariant of $S = \Sigma \times \mathbb{P}^1$ with $w = c_1$ and polarisation H .

Proof. By definition, the right hand side is $\epsilon_S(c_1) < \alpha^a \beta^b \psi_{i_1} \cdots \psi_{i_r}, [\mathfrak{M}] >$, where $\alpha = 2\mu(\Sigma) \in H^2(\mathfrak{M})$, $\beta = -4\mu(x) \in H^4(\mathfrak{M})$, $\psi_i = \mu(\gamma_i^\#) \in H^3(\mathfrak{M})$. Here the factor $\epsilon_S(c_1) = (-1)^{\frac{\kappa_S c_1 + c_1^2}{2}} = (-1)^{g-1}$ compares the complex orientation of $\mathfrak{M}$ and its natural orientation as a moduli space of anti-self-dual connections [4]. By [9, theorem 10], this is worked out to be $(-1)^{g-1} < (4\omega + X)^a (X^2)^b \phi_{i_1} \cdots \phi_{i_r} X^r, [J] >$, where $X^{2g-1+i} = s_i(\mathcal{E}_\zeta) = \frac{(-8)^i}{i!} \omega^i$. Thus the theorem follows from lemmas 8 and 10. $\square$

Remark 12. The formula in theorem 11 is not right for $g = 2$, as in such case, the quantum product $h^2 \in QH^*(N)$ differs from β by a quantum correction.

Remark 13. Suppose $g \geq 2$ and let $\alpha^a \beta^b \psi_{i_1} \cdots \psi_{i_r} \in \mathbb{A}(\Sigma)$ have degree $6g - 6$. Then

$$\begin{aligned} \Psi_0^{M_\Sigma}(\alpha, \cdot^{(a)}, \alpha, \beta, \cdot^{(b)}, \beta, \psi_{i_1}, \dots, \psi_{i_r}) &= \epsilon_S(\mathbb{P}^1) < \alpha^a \beta^b \psi_{i_1} \cdots \psi_{i_r}, [M_\Sigma] > \\ &= -D_{S,H}^{\mathbb{P}^1}((2\Sigma)^a (-4pt)^b \gamma_{i_1}^\# \cdots \gamma_{i_r}^\#), \end{aligned}$$

as the moduli space of anti-self-dual connections on S of dimension $6g - 6$ is M_Σ .

5. QUANTUM COHOMOLOGY RING OF M_Σ

The action of the mapping class group on M_Σ is symplectic, so the quantum product restricts to the invariant part of the cohomology, thus having defined $QH_I^*(M_\Sigma)$ (see [16, section 3.1]). To give a description of it, let us define

$$(11) \quad \begin{cases} \hat{\alpha} = \alpha \\ \hat{\beta} = \beta + r_g \\ \hat{\gamma} = -2 \sum \psi_i \psi_{i+g} \end{cases}$$

where $r_g \in \mathbb{Q}$, $g \geq 1$ (to be determined shortly), and $\hat{\gamma}$ is given using the quantum product. It might happen that $\hat{\gamma} = \gamma + s_g \alpha$, $s_g \in \mathbb{Q}$. The need of introducing this quantum corrections in the generators was noticed already in [1].

As a consequence of the description of $H_I^*(M_\Sigma)$ given in section 2, the results in [18] imply that there is a presentation

$$QH_I^*(M_\Sigma) = \mathbb{Q}[\hat{\alpha}, \hat{\beta}, \hat{\gamma}] / J_g,$$

where the ideal J_g is generated by three elements Q_g^1 , Q_g^2 and Q_g^3 which are deformations graded mod 4 of q_g^1 , q_g^2 and q_g^3 , respectively (see [16]). This means that $J_g = (Q_g^1, Q_g^2, Q_g^3)$ for

$$(12) \quad Q_g^i = \sum_{j \geq 0} Q_{g,j}^i,$$

where $\deg(Q_{g,j}^i) = \deg(q_g^i) - 4j$, $j \geq 0$, and $Q_{g,0}^i = q_g^i$. There is still one source of possible ambiguity coming from adding a scalar multiple of Q_g^1 to Q_g^3 . To avoid this, we require the coefficient of $\hat{\alpha}^g$ in Q_g^3 to be zero. Recall the main result from [10].

Theorem 14 ([10]). *Define $R_0^1 = 1$, $R_0^2 = 0$, $R_0^3 = 0$ and then recursively, for all $g \geq 1$,*

$$\begin{cases} R_{g+1}^1 = \alpha R_g^1 + g^2 R_g^2 \\ R_{g+1}^2 = (\beta + (-1)^{g+1} 8) R_g^1 + \frac{2g}{g+1} R_g^3 \\ R_{g+1}^3 = \gamma R_g^1 \end{cases}$$

Then the invariant part of the instanton Floer cohomology of $\Sigma \times \mathbb{S}^1$ is $HF^(\Sigma \times \mathbb{S}^1)_I = \mathbb{Q}[\alpha, \beta, \gamma]/(R_g^1, R_g^2, R_g^3)$. R_g^1 , R_g^2 , R_g^3 are uniquely determined by the conditions that the leading term of R_g^i is q_g^i , $i = 1, 2, 3$ and that the coefficient of α^g in R_g^3 is zero.*

Accounting for the difference in signs between theorem 11 and remark 13, the prospective generators of J_g are defined as follows.

Definition 15. For $i = 1, 2, 3$ and $g \geq 1$, set

$$\hat{R}_g^i(\hat{\alpha}, \hat{\beta}, \hat{\gamma}) = (\sqrt{-1})^{-\deg q_g^i} R_g^i(\sqrt{-1}^g \hat{\alpha}, \sqrt{-1}^{2g} \hat{\beta}, \sqrt{-1}^{3g} \hat{\gamma}),$$

i.e. when g is even, $\hat{R}_g^i = R_g^i$ and when g is odd, $\hat{R}_g^i$ is obtained from R_g^i by changing the sign of the homogeneous components of degrees $\deg(q_g^i) - 4 - 8j$, $j \geq 0$.

Thus we expect that $Q_g^i = \hat{R}_g^i$, $i = 1, 2, 3$ (i.e. $J_g = (\hat{R}_g^1, \hat{R}_g^2, \hat{R}_g^3)$), for $g \geq 1$ (see [1] [16]). Let us review the known cases.

Example 16. For $g = 1$, we set $r_1 = -8$, so that $\hat{\alpha} = \alpha$, $\hat{\beta} = \beta - 8$, $\hat{\gamma} = \gamma$. Then the ideal of relations is generated by $\hat{R}_1^1 = \hat{\alpha}$, $\hat{R}_1^2 = \hat{\beta} + 8$ and $\hat{R}_1^3 = \hat{\gamma}$. The correction $r_1 = -8$ is arranged in such a way that things work.

Example 17. For $g = 2$, the quantum cohomology ring $QH^*(M_\Sigma)$ has been computed by Donaldson [3], using an explicit description of M_Σ as the intersection of two quadrics in $\mathbb{P}^5$. Let h_2 , h_4 and h_6 be the integral generators of $QH^2(M_\Sigma)$, $QH^4(M_\Sigma)$ and $QH^6(M_\Sigma)$, respectively. Then, with our notations, $\alpha = h_2$, $\beta = -4h_4$ and $\gamma = 4h_6$ (see [1]). The computations in [3] yield $\hat{\gamma} = \gamma - 4\alpha$. Now we set $r_2 = 4$, i.e. $\hat{\beta} = \beta + 4$. It is now easy to check that the relations found in [3] can be translated to the relations $\hat{R}_2^1 = \hat{\alpha}^2 + \hat{\beta} - 8$, $\hat{R}_2^2 = (\hat{\beta} + 8)\hat{\alpha} + \hat{\gamma}$ and $\hat{R}_2^3 = \hat{\alpha}\hat{\gamma}$ for $QH_I^*(M_\Sigma)$.

The artificially introduced term $r_2 = 4$ is due to the same phenomenon which causes the failure of lemma 10 for $g = 2$, i.e. the quantum product h^2 differs from β in (9) (defined with the cup product) because of a quantum correction in $QH^*(N)$ which appears when $g = 2$.

In the general case we have

Theorem 18. *Let $g \geq 3$. Put $r_g = 0$, so that $\hat{\alpha} = \alpha$ and $\hat{\beta} = \beta$. Then we can write*

$$\begin{cases} Q_g^1 = \hat{R}_g^1 + f_g^1 \\ Q_g^2 = \hat{R}_g^2 + f_g^2 \\ Q_g^3 = \hat{R}_g^3 + f_g^3 \end{cases}$$

where $\deg(f_g^i) \leq \deg(q_g^i) - 8$, $i = 1, 2, 3$.

Proof. As in equation (12), we can write $\hat{R}_g^i = \sum_{j \geq 0} \hat{R}_{g,j}^i$, where $\deg(\hat{R}_{g,j}^i) = \deg(q_g^i) - 4j$, $j \geq 0$ (and analogously for R_g^i). Clearly, $\hat{R}_{g,0}^i = q_g^i = Q_{g,0}^i$, for $i = 1, 2, 3$. We want to check that $Q_{g,1}^i = \hat{R}_{g,1}^i = (-1)^g R_{g,1}^i$. Pick any $z = \alpha^a \beta^b \psi_{i_1} \cdots \psi_{i_r} \in \mathbb{A}(\Sigma)$ of degree $6g - 2 - \deg(q_g^i)$. By theorem 14, $\phi^w(\Sigma \times D^2, R_g^i) = 0$ (see [10] for notations), so $D_{S,H}^{(w,\Sigma)}(R_g^i z) = 0$, i.e.

$$D_{S,H}^{\mathbb{P}^1}(R_{g,1}^i z) + D_{S,H}^{c_1}(R_{g,0}^i z) = 0.$$

From theorem 11 and remark 13 this is translated as the component in $H^{6g-6}(M_\Sigma)$ of the quantum product $-R_{g,1}^i z + (-1)^{g-1} R_{g,0}^i z \in QH^*(M_\Sigma)$ vanishing. On the other hand, by definition $Q_g^i z = 0 \in QH^*(M_\Sigma)$, so the component in $H^{6g-6}(M_\Sigma)$ of the quantum product $Q_{g,1}^i z + Q_{g,0}^i z \in QH^*(M_\Sigma)$ is zero. Thus $\langle Q_{g,1}^i, z \rangle = \langle (-1)^g R_{g,1}^i, z \rangle$, for any z of degree $6g - 2 - \deg(q_g^i)$, and hence $Q_{g,1}^i \equiv (-1)^g R_{g,1}^i \pmod{I_g}$ (I_g is the ideal defined in section 2). This gives the required equality $Q_{g,1}^i = \hat{R}_{g,1}^i$ (in the case $i = 3$ we have to use the vanishing of the coefficient of $\hat{\alpha}^g$ for both Q_g^3 and R_g^3). $\square$

6. THE CASE OF GENUS $g = 3$

It is natural to ask to what extent the first quantum correction determine the full structure of the quantum cohomology of M_Σ . In [3], Donaldson finds the first quantum correction for M_Σ when the genus of Σ is $g = 2$ and proves that this determines the quantum product. In this section we are going to check that this also happens for $g = 3$, finding thus the quantum cohomology of the moduli space of stable bundles over a Riemann surface of genus $g = 3$.

Proposition 19. *Let Σ have genus $g = 3$. Then $Q_3^1 = \hat{R}_3^1$, $Q_3^2 = \hat{R}_3^2$ and $Q_3^3 = \hat{R}_3^3$, i.e.*

$$QH_I^*(M_\Sigma) = \mathbb{Q}[\hat{\alpha}, \hat{\beta}, \hat{\gamma}] / (\hat{R}_3^1, \hat{R}_3^2, \hat{R}_3^3).$$

Proof. Theorem 18 says that

$$\begin{cases} Q_g^1 = \hat{R}_g^1 = \hat{\alpha}(\hat{\alpha}^2 + \hat{\beta} + 8) + 4(\hat{\alpha}\hat{\beta} - 8\hat{\alpha} + \hat{\gamma}) \\ Q_g^2 = \hat{R}_g^2 + x = (\hat{\beta} + 8)(\hat{\alpha}^2 + \hat{\beta} + 8) + \frac{4}{3}\hat{\alpha}\hat{\gamma} + x \\ Q_g^3 = \hat{R}_g^3 + y\hat{\alpha} = \hat{\gamma}(\hat{\alpha}^2 + \hat{\beta} + 8) + y\hat{\alpha} \end{cases}$$

where $x, y \in \mathbb{Q}$. The main tool that we shall use is the nilpotency of $\hat{\gamma}$. Actually, from its definition (11), $\hat{\gamma}^4 = 0$ in $QH^*(M_\Sigma)$.

Suppose $y \neq 0$. Then the third relation implies $\hat{\alpha}^4 = 0$. Multiplying the first relation by $\hat{\alpha}$ we get $5\hat{\alpha}^2\hat{\beta} + 24\hat{\alpha}^2 + 4\hat{\alpha}\hat{\gamma} = 0$. This is a relation of degree $2g + 2 = 8$. Thus it must be a multiple of Q_g^2 , which is not. This contradiction implies $y = 0$.

Let us see $x = 0$. It is easy to check that

$$\hat{\gamma}^3 = \frac{\hat{\gamma}^2}{4}\hat{R}_3^1 - \frac{3\hat{\gamma}(\hat{\beta} - 8)}{4}\hat{R}_3^2 + \frac{3(\hat{\beta} + 8)(\hat{\beta} - 8) - \hat{\alpha}\hat{\gamma}}{4}\hat{R}_3^3.$$

The relations above imply that $\hat{\gamma}^3 = -\frac{3}{4}\hat{\gamma}(\hat{\beta} - 8)x$ in $QH^*(M_\Sigma)$. So $\hat{\gamma}^2(\hat{\beta} - 8) = 0$. Also, the third relation (with $y = 0$) gives $\hat{\gamma}\hat{\alpha}^2 = -\hat{\gamma}(\hat{\beta} + 8)$. Thus $\hat{\gamma}^2\hat{\alpha}^2 = -16\hat{\gamma}^2$ and $\hat{\gamma}^2\hat{\beta} = 8\hat{\gamma}^2$. Multiply the first relation by $\hat{\gamma}^2$ to get $\hat{\gamma}^3 = 0$. As $\hat{\gamma}(\hat{\beta} - 8) \neq 0$ (because this is not a multiple of the only relation of degree $2g = 6$), it must be $x = 0$. $\square$

Corollary 20. *Let Σ be a Riemann surface of genus $g = 3$. Then*

$$QH^*(M_\Sigma) = \bigoplus_{k=0}^{g-1} \Lambda_0^k H^3 \otimes \mathbb{Q}[\hat{\alpha}, \hat{\beta}, \hat{\gamma}] / \hat{I}_{g-k}$$

where we put $\hat{I}_r = (R_r^1(\sqrt{-1}^g \hat{\alpha}, \sqrt{-1}^{2g} \hat{\beta}, \sqrt{-1}^{3g} \hat{\gamma}), R_r^2(\sqrt{-1}^g \hat{\alpha}, \sqrt{-1}^{2g} \hat{\beta}, \sqrt{-1}^{3g} \hat{\gamma}), R_r^3(\sqrt{-1}^g \hat{\alpha}, \sqrt{-1}^{2g} \hat{\beta}, \sqrt{-1}^{3g} \hat{\gamma})), 1 \leq r \leq g$.

Proof. This is an easy consequence of the former proposition and [10, lemma 7], noting that the hypothesis only has to be checked up to $g = 3$. One has to be careful with the exponents of $\sqrt{-1}$ everywhere. $\square$

REFERENCES

1. M. Bershadsky, A. Johansen, V. Sadov and C. Vafa, Topological reduction of 4D SYM to 2D σ -models, Preprint, 1995.
2. U. Desale and S. Ramanan, Classification of vector bundles of rank 2 on hyperelliptic curves, *Inventiones Math.* **38** 1976, 161-185.

3. S. K. Donaldson, Floer homology and algebraic geometry, *Vector bundles in algebraic geometry*, London Math. Soc. Lecture Notes Series, **208** Cambridge University Press, Cambridge, 1995, 119-138.
4. S. K. Donaldson and P. B. Kronheimer, *The geometry of 4-manifolds*, Oxford University Press, 1990.
5. G. Ellingsrud and L. Göttsche, Variation of moduli spaces and Donaldson invariants under change of polarisation, *Journal reine angew. Math.* **467** 1995, 1-49.
6. A. D. King and P. E. Newstead, On the cohomology ring of the moduli space of rank 2 vector bundles on a curve, Liverpool Preprint, 1994.
7. P. B. Kronheimer and T. S. Mrowka, Embedded surfaces and the structure of Donaldson's polynomial invariants, *Jour. Differential Geometry*, **41** 1995, 573-734.
8. D. McDuff and D. A. Salamon, J-holomorphic curves and quantum cohomology, Preprint.
9. V. Muñoz, Wall-crossing formulae for algebraic surfaces with $q > 0$, alg-geom/9709002.
10. V. Muñoz, Ring structure of the Floer cohomology of $\Sigma \times \mathbb{S}^1$, dg-ga/9710029.
11. S. Piunikhin, Quantum and Floer cohomology have the same ring structure, MIT Preprint, 1994.
12. S. Piunikhin, D. Salamon and M. Schwarz, Symplectic Floer-Donaldson theory and quantum cohomology, Warwick Preprint, 1995.
13. Z. Qin, Moduli of stable sheaves on ruled surfaces and their Picard groups, *Jour. Reine angew. Math.* **433** 1992, 201-219.
14. Y. Ruan, Topological sigma model and Donaldson type invariants in Gromov theory, *Duke Math. Jour.* **83** 1996, 461-500.
15. Y. Ruan and G. Tian, A mathematical theory of quantum cohomology, *Jour. Diff. Geom.* **42** 1995, 259-367.
16. B. Siebert, An update on (small) quantum cohomology, Preprint, 1997.
17. B. Siebert and G. Tian, Recursive relations for the cohomology ring of moduli spaces of stable bundles, *Proceedings of 3rd Gökova Geometry-Topology Conference 1994*.
18. B. Siebert and G. Tian, On quantum cohomology rings of Fano manifolds and a formula of Vafa and Intriligator, submitted to Duke Math. Journal.
19. M. Thaddeus, Conformal field theory and the cohomology of the moduli space of stable bundles, *Jour. Differential Geometry*, **35** 1992, 131-150.

DEPARTAMENTO DE ÁLGEBRA, GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS, UNIVERSIDAD DE MÁLAGA, 29071 MÁLAGA, SPAIN

E-mail address: vmunoz@agt.cie.uma.es